

4-1992

# Some Facts About Cycels and Tidy Groups

Kevin O'Bryant

*Rose-Hulman Institute of Technology*

D. Patrick

*Carnegie Mellon University*

Lawren Smithline

*Harvard University*

Eric Wepsic

*Harvard University*

Advisors:

Gary Sherman

Follow this and additional works at: [http://scholar.rose-hulman.edu/math\\_mstr](http://scholar.rose-hulman.edu/math_mstr)

 Part of the [Algebra Commons](#)

---

## Recommended Citation

O'Bryant, Kevin; Patrick, D.; Smithline, Lawren; and Wepsic, Eric, "Some Facts About Cycels and Tidy Groups" (1992). *Mathematical Sciences Technical Reports (MSTR)*. 131.

[http://scholar.rose-hulman.edu/math\\_mstr/131](http://scholar.rose-hulman.edu/math_mstr/131)

MSTR 92-04

This Article is brought to you for free and open access by the Mathematics at Rose-Hulman Scholar. It has been accepted for inclusion in Mathematical Sciences Technical Reports (MSTR) by an authorized administrator of Rose-Hulman Scholar. For more information, please contact [weir1@rose-hulman.edu](mailto:weir1@rose-hulman.edu).

**SOME FACTS ABOUT CYCELS  
AND TIDY GROUPS**

**K. O'Bryant, D. Patrick, L. Smithline and E. Wepsic**

**MS TR 92-04**

**April 1992**

**Department of Mathematics  
Rose-Hulman Institute of Technology  
Terre Haute, IN 47803**

**FAX(812) 877-3198**

**Phone: (812) 877-8391**

---

# Some Facts About Cycles and Tidy Groups

K. O'Bryant\*    D. Patrick\*    L. Smithline\*    E. Wepsic\*

## 1 Definitions

Note: In this paper, all groups mentioned are finite. We denote the identity element of a group  $G$  by  $e$ , and  $G \setminus \{e\}$  by  $G^\times$ . Also,  $H \leq G$  denotes that  $H$  is a subgroup of  $G$ .

Recall that the centralizer of an element  $x \in G$  can be defined by

$$C(x) = \{y \in G \mid \langle x, y \rangle \text{ is abelian}\}.$$

If, in the above definition, we replace the word “abelian” with the word “cyclic”, we get a subset of the centralizer, called the *cyclicizer*. To be explicit, define the cyclicizer of an element  $x \in G$ , denoted  $\text{Cyc}(x)$ , by

$$\text{Cyc}(x) = \{y \in G \mid \langle x, y \rangle \text{ is cyclic}\}.$$

Some properties of cyclicizers are discussed in [1].

Also, just as the center of the group can be defined by

$$Z(G) = \bigcap_{x \in G} C(x),$$

we may define a similar construct, called the *cyce*<sup>1</sup>, denoted  $K(G)$ , by

$$K(G) = \bigcap_{x \in G} \text{Cyc}(x).$$

---

\*Supported by NSF grant number DMS-910059.

<sup>1</sup>The notation for the cycle comes from the Hungarian word *kerek*, which means “round”.

## 2 Properties of the cycl

### 2.1 Basic properties

It is clear from the definitions above that  $Cyc(x) \subseteq C(x)$  for all  $x \in G$ , and thus  $K(G) \subseteq Z(G)$ . It is also clear that for all  $g \in K(G)$  and all  $x \in G$ , we have  $\langle x, g \rangle$  cyclic. This allows us to prove the following:

**Theorem 1** For all groups  $G$ ,  $K(G) \triangleleft G$ .

*Proof:* Let  $g, h \in K(G)$  and  $x \in G$  be given. Then  $\langle g^{-1}, x \rangle = \langle g, x \rangle$ , which is cyclic, so  $g^{-1} \in K(G)$ . Also,  $\langle gh, x \rangle \leq \langle g, h, x \rangle$ , which is cyclic, so  $gh \in K(G)$ . So  $K(G) \leq G$ . But  $K(G) \leq Z(G)$  as well, so  $K(G)$  is normal. ■

Recall that  $Z(G)$  can also be defined as the intersection of the maximal abelian subgroups of  $G$ . We can characterize  $K(G)$  similarly.

**Theorem 2** For all groups  $G$ ,  $K(G)$  is the intersection of the maximal cyclic subgroups of  $G$ .

*Proof:* Let  $x$  be in the intersection of the maximal cyclic subgroups of  $G$ , and pick arbitrary  $g \in G$ . Then  $g$  is contained in some maximal cyclic subgroup  $\langle h \rangle$  of  $G$ . But  $x \in \langle h \rangle$  as well, so  $\langle x, g \rangle \leq \langle h \rangle$ . Hence  $g$  and  $x$  generate a cyclic subgroup. Since this is true for all  $g \in G$ , we have that  $x \in K(G)$ .

Conversely, Let  $x \in K(G)$  and  $g \in G$  be given such that  $\langle g \rangle$  is a maximal cyclic subgroup of  $G$ . Then  $\langle x, g \rangle$  is cyclic, but since  $\langle g \rangle$  is maximal we must have  $\langle x, g \rangle = \langle g \rangle$ . Thus  $x \in \langle g \rangle$ . Since this is true for all maximal cyclic subgroups of  $G$ , we must have  $x$  in the intersection of the maximal cyclic subgroups of  $G$ . ■

An immediate corollary of Theorem 2 is

**Corollary 3** For all groups  $G$ ,  $K(G)$  is cyclic. □

Let us summarize some of the similarities of the center and the cycl:

$Z(G)$	$K(G)$
Intersection of centralizers	Intersection of cyclicizers
Intersection of maximal abelian subgroups	Intersection of maximal cyclic subgroups
$\langle x, g \rangle$ abelian for $x \in Z(G), g \in G$	$\langle x, g \rangle$ cyclic for $x \in K(G), g \in G$
$Z(G)$ abelian	$K(G)$ cyclic

We may also wish to consider the cyclic analog of nilpotency, which could be termed “cycelpotency”, by constructing the “ascending cycl series”:

$$\langle e \rangle, K(G), K(G/K(G)), \dots$$

However, this concept proves to be trivial, as shown in the following theorem:

**Theorem 4** For all groups  $G$ ,  $K(G/K(G)) = \langle e \rangle$ .

*Proof:* Let  $K = K(G) = \langle k \rangle$ , by Corollary 3. Suppose  $L \in K(G/K(G))$ . For all  $c \in G$ , where  $C = cK$ , there is a  $D \in G/K$  such that  $\langle L, C \rangle = \langle D \rangle$ . But  $L = lK, D = dK$  for some  $l, d \in G$ , and hence

$$\langle l, c, k \rangle = \langle l, c \rangle K = \langle d \rangle K = \langle d, k \rangle = \langle d' \rangle$$

for some  $d' \in G$ . In particular  $\langle l, c, k \rangle$  is cyclic, thus  $\langle l, c \rangle \leq \langle l, c, k \rangle$  is cyclic. But  $c \in G$  is arbitrary, so  $l \in K(G)$  and hence  $L = e$ . ■

## 2.2 Miscellaneous properties

The above theorem deals with the cycl of  $G/K(G)$ . We can also prove a similar result about  $G/Z(G)$ . First we need the following technical lemma.

**Lemma 5** For all groups  $G$ , if  $gZ \in (G/Z(G))^*$ , then  $\langle \text{Cyc}(gZ) \rangle \neq G/Z(G)$ .

*Proof:* Suppose  $hZ \in \text{Cyc}(gZ)$ . Then  $\langle gZ, hZ \rangle$  is cyclic in  $G/Z(G)$ , so  $\langle g, h, Z \rangle$  is abelian in  $G$ . Thus  $h \in C(g)$ . Thus, since  $C(g) \leq G$ , we have that if  $h$  is in the preimage of  $\langle \text{Cyc}(gZ) \rangle$ , then  $h \in C(g)$ . Thus, since  $gZ \neq e$ , we have that  $C(g) \neq G$ , so  $\langle \text{Cyc}(gZ) \rangle \neq G/Z$ . ■

This immediately gives the following:

**Theorem 6** For all groups  $G$ ,  $K(G/Z(G)) = \langle e \rangle$ .

*Proof:* This follows immediately from Lemma 5, since if  $\bar{g} \in K(G/Z(G))$ , then  $\text{Cyc}(\bar{g}) = G/Z$ , so by Lemma 5 we must have  $\bar{g} = e$ . ■

Lemma 5 also gives us the following interesting result on the structure of  $G/Z$ . Recall that a non-abelian group is said to be *Dedekind* if all of its proper subgroups are normal.

**Theorem 7** For all groups  $G$ ,  $G/Z$  is not a Dedekind group.

*Proof:* Suppose  $G/Z$  is a Dedekind group. Then  $G/Z = Q \times A$ , where  $Q$  is the quaternion group and  $A$  is an abelian group. Consider the element  $(-1, 0)$  of  $G/Z$ . Then  $\text{Cyc}((-1, 0)) = (Q \times \{0\}) \cup (Q \setminus \{-1, 1\} \times A)$ , and thus  $\langle \text{Cyc}((-1, 0)) \rangle = G/Z$ , a contradiction of Lemma 5. ■

We can also characterize the  $p$ -groups with non-trivial cycles.

**Theorem 8** Suppose  $G$  is a  $p$ -group. Then  $K(G) \neq \langle e \rangle$  if and only if  $G$  is cyclic or generalized quaternion. Moreover, if  $K(G) \neq \langle e \rangle$ , then  $K(G) = Z(G)$ .

*Proof:* If  $G$  is cyclic, then  $K(G) = Z(G) = G$ . If  $G$  is generalized quaternion, then every maximal cyclic subgroup contains the center (see [2]), so by Theorem 2,  $K(G) = Z(G)$ .

Conversely, suppose that  $G$  is a  $p$ -group with non-trivial cycle. Then by Corollary 3,  $K(G)$  is cyclic, so  $K(G)$  has exactly  $p - 1$  elements of order  $p$ . Suppose there is another element  $x \in G$ , with  $x \notin K(G)$ , such that  $|x| = p$ . Consider  $H$ , the maximal cyclic subgroups containing  $x$ . Then by the previous lemma,  $K(G) \leq H$ . But  $H$  is cyclic, and contains  $p$  elements of order  $p$ , a contradiction. Hence,  $G$  has exactly  $p - 1$  elements of order  $p$ , and by a well-known result (in [2], among other sources),  $G$  must be either cyclic or generalized quaternion. ■

Finally, we have the result which indicates when a non-cyclic group  $G$  is spanned by a minimal number of cyclic subgroups.

**Theorem 9**  $G$  is the union of three proper cyclic subgroups if and only if  $G$  is isomorphic to either  $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{C}$  or  $\mathbf{Q} \times \mathbf{C}$ , where  $\mathbf{C}$  is a cyclic group of odd order, and  $\mathbf{Q}$  is the quaternion group of order 8.

*Proof:* It is clear that both  $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{C}$  and  $\mathbf{Q} \times \mathbf{C}$  are the union of three proper cyclic subgroups. Bruckheimer, Bryan and Muir [3] have shown that a group is the union of three proper subgroups if and only if their intersection  $N$  is normal and  $G/N \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ . It follows from Theorem 1, Theorem 2, and [3] that the groups spanned by three cyclic subgroups are precisely those for which  $G/K(G) \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ . We now classify such groups.

First, note that  $G$  is nilpotent. This can be seen by observing that

$$G/Z(G) \cong (G/K(G))/(Z(G)/K(G))$$

is a factor group of an abelian group, and so is abelian. Thus, we can write  $G \cong Syl_2 \times Syl_3 \times Syl_5 \times \dots$ . Since  $|G/K| = 2^2$ ,  $K(G)$  must contain  $Syl_3 \times Syl_5 \times \dots$ . It follows from Corollary 3 that  $C \cong Syl_3 \times Syl_5 \times \dots$  is cyclic and of odd order. Note that we have also proven that  $|K(G)| = 2^a|C|$ , where  $a$  is a nonnegative integer.

Suppose  $K(G) \cap Syl_2 = \langle e \rangle$ . Then  $|Syl_2| = |G/K(G)| = 4$ . If  $Syl_2$  were cyclic, then  $G$  would be as well. In that case, however,  $G$  would not be the union of three proper cyclic subgroups. Hence  $Syl_2 \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ , and  $G \cong \mathbf{Z}_2 \times \mathbf{Z}_2 \times C$ . Now, suppose that  $K(G) \cap Syl_2 \neq \langle e \rangle$ . It is clear that  $K(G) \cap Syl_2 \leq K(Syl_2)$ , from which we see that  $Syl_2$  is a 2-group with a non-trivial center. It follows from Theorem 8 that  $Syl_2$  is cyclic or generalized quaternion. If  $Syl_2$  were cyclic, then  $G$  would also be cyclic and, consequently, not the union of three proper cyclic subgroups. If  $Syl_2$  is generalized quaternion, then by Theorem 8,  $|K(Syl_2)| = |Z(Syl_2)|$ . But the center of a generalized quaternion is of order 2 (see [2]). We can now see that  $|K(G) \cap Syl_2| \leq |K(Syl_2)| = 2$ , and so  $|K(G)| = 2^a|C| = 2|C|$ . Thus,  $|G| = |G/K(G)| \cdot |K(G)| = 4 \cdot 2|C|$ . Hence,  $|Syl_2| = 8$  and  $Syl_2$  is the quaternion group of order 8. ■

### 3 Tidy groups

Note that unlike its analog the centralizer, a cyclicizer is not necessarily a subgroup. For example, in the group  $\mathbf{Z}_4 \times \mathbf{Z}_2$ , the cyclicizer of the element  $(2, 0)$  is of order 6:

$$\text{Cyc}((2, 0)) = \{(0, 0), (1, 0), (1, 1), (2, 0), (3, 0), (3, 1)\}.$$

A natural question to ask is: what can be said about those groups in which all of the cyclicizers are actually subgroups?

Let  $G$  be a group.  $G$  is said to be *tidy* if  $\text{Cyc}(x)$  is a subgroup for all  $x \in G$ . We begin with the following lemma on direct products of tidy groups.

**Lemma 10** *Let  $G$  and  $H$  be groups such that  $|G|$  and  $|H|$  are relatively prime. Then  $G \times H$  is tidy if and only if  $G$  and  $H$  are tidy.*

*Proof:* Suppose  $G$  and  $H$  are such that  $|G|$  and  $|H|$  are relatively prime, and that  $G$  and  $H$  are both tidy. Then for all  $(g, h) \in G \times H$ ,  $\text{Cyc}((g, h)) = \text{Cyc}(g) \times \text{Cyc}(h)$  (since for all  $(g_1, h_1), (g_2, h_2) \in G \times H$ , we have  $\langle (g_1, h_1), (g_2, h_2) \rangle = \langle g_1, g_2 \rangle \times \langle h_1, h_2 \rangle$ ). Hence  $\text{Cyc}((g, h))$  is a subgroup.

Conversely, suppose that  $G \times H$  is tidy. Then for all  $g \in G$ , we have that  $\text{Cyc}(g) = \pi_G(\text{Cyc}((g, e)))$  (where  $\pi_G$  is the projection homomorphism from  $G \times H$  to  $G$ ). So  $\text{Cyc}(g)$  is a subgroup. ■

We now classify all the abelian tidy groups.

**Theorem 11** *Let  $G$  be an abelian group. Then  $G$  is tidy if and only if every  $p$ -Sylow subgroup of  $G$  is cyclic or elementary abelian.*

*Proof:* By the previous lemma,  $G$  is tidy if and only if each  $p$ -Sylow subgroup of  $G$  is tidy. Let  $P$  be a  $p$ -Sylow subgroup of  $G$ . If  $P$  is cyclic, then for all  $x \in P$ , we have  $\text{Cyc}(x) = P$ . If  $P$  is elementary abelian, then every  $x \in P^\times$  is of order  $p$ , so  $\text{Cyc}(x) = \langle x \rangle$ . Otherwise,  $P$  contains a subgroup isomorphic to  $\mathbb{Z}_p \oplus \mathbb{Z}_{p^2}$ , which is not tidy (since the element  $(0, p)$  has a cyclicizer which is not a subgroup). ■

We also classify tidy  $p$ -groups. First we need the following lemma:

**Lemma 12** *Suppose  $G$  is a tidy  $p$ -group. Let  $x \in G$  be such that  $|x| \neq p$ , and  $\langle x \rangle$  is a maximal cyclic subgroup. Then  $Z(G) \leq \langle x \rangle$ .*

*Proof:* Let  $z \in Z(G)$  be given, and consider  $x$  as above. Then  $\langle z, x \rangle$  is abelian. Furthermore  $\langle z, x \rangle$  is not elementary abelian, since  $|x| \neq p$ . So by Theorem 11,  $\langle z, x \rangle$  is cyclic. But  $\langle x \rangle$  is maximal; hence  $z \in \langle x \rangle$ . Since  $z \in Z(G)$  was arbitrary, we conclude that  $Z(G) \leq \langle x \rangle$ . ■

An immediate corollary is

**Corollary 13** *Suppose  $G$  is a tidy  $p$ -group. Then  $Z(G)$  is cyclic. □*

This allows us to prove the following:

**Theorem 14** *Suppose  $G$  is a  $p$ -group. Then  $G$  is tidy if and only if there exists  $H \triangleleft G$ , where  $H$  is cyclic or generalized quaternion, and for all  $x \in G \setminus H$ , we have  $|x| = p$ .*

*Proof:* Suppose  $G$  is a tidy  $p$ -group. Let  $H = \text{Cyc}(z)$ , where  $z$  is a non-identity element of  $Z(G)$ . First, consider any element  $x \in G \setminus H$ , with  $|x| \neq p$ . Then  $x$  is contained in some maximal cyclic subgroup  $\langle y \rangle$ , with  $|y| \neq p$ , so by Lemma 12,  $z \in \langle y \rangle$ . But then  $\langle x, z \rangle \in \langle y \rangle$ , so  $\langle x, z \rangle$  is cyclic, and hence  $x \in \text{Cyc}(z) = H$ .

Next, we observe that  $z \in K(H)$ , since  $\langle z, h \rangle$  is cyclic for any  $h \in H$ , so by Theorem 8,  $H$  is either cyclic or generalized quaternion. Finally, if  $\langle h \rangle$  is



a maximal cyclic subgroup of  $H$ , with  $|h| > p$ , then any conjugate of  $\langle h \rangle$  is a cyclic subgroup of order greater than  $p$ , and hence must be in  $H$ . So  $H \triangleleft G$ .

For the converse, suppose  $G$  is a  $p$ -group with a normal subgroup  $H$  as above. Then if  $x \in H$ ,  $\text{Cyc}(x)$  is a subgroup (since cyclic and generalized quaternion groups are themselves tidy), and if  $x \notin H$ , then  $|x| = p$ , so  $\text{Cyc}(x) = \langle x \rangle$ . So  $\text{Cyc}(x)$  is a subgroup for all  $x \in G$ , and hence  $G$  is tidy. ■

## 4 References

- [1] D. Patrick and E. Wepsic, "Cyclicizers, Centralizers, and Normalizers", Rose-Hulman Technical Report MS-TR 91-05.
- [2] D. Gorenstein, *Finite Groups*, New York: Harper & Row, 1968.
- [3] M. Bruckheimer, A.C. Bryan, and A. Muir, "Groups Which Are the Union of Three Subgroups", *American Mathematical Monthly* 77 (1970).

David Patrick  
Carnegie Mellon Univ.  
Pittsburgh, PA 15213

Eric Wepsic & Lawren Smithline  
Harvard University  
Cambridge, MA 02138

Kevin O'Bryant  
Rose-Hulman Inst. of Tech.  
Terre Haute, IN 47803