

The Log Convex Density Conjecture in Hyperbolic Space

Leonardo Di Giosia
Lewis and Clark College

Jahangir Habib
Williams College

Lea Kenigsberg
Stony Brook University

Dylanger Pittman
Williams College

Weitao Zhu
Williams College

Follow this and additional works at: <https://scholar.rose-hulman.edu/rhumj>

Recommended Citation

Di Giosia, Leonardo; Habib, Jahangir; Kenigsberg, Lea; Pittman, Dylanger; and Zhu, Weitao (2017) "The Log Convex Density Conjecture in Hyperbolic Space," *Rose-Hulman Undergraduate Mathematics Journal*: Vol. 18 : Iss. 1 , Article 9.

Available at: <https://scholar.rose-hulman.edu/rhumj/vol18/iss1/9>

ROSE-
HULMAN
UNDERGRADUATE
MATHEMATICS
JOURNAL

THE LOG CONVEX DENSITY
CONJECTURE IN HYPERBOLIC SPACE

Leonardo Di Giosia^a Jahangir Habib^b
Lea Kenigsberg^c Dylanger Pittman^d Weitao Zhu^e

VOLUME 18, No. 1, SPRING 2017

Sponsored by

Rose-Hulman Institute of Technology
Department of Mathematics
Terre Haute, IN 47803
mathjournal@rose-hulman.edu
scholar.rose-hulman.edu/rhumj

^aLewis & Clark College

^bWilliams College

^cStony Brook University

^dWilliams College

^eWilliams College

ROSE-HULMAN UNDERGRADUATE MATHEMATICS JOURNAL
VOLUME 18, No. 1, SPRING 2017

THE LOG CONVEX DENSITY CONJECTURE IN
HYPERBOLIC SPACE

Leonardo Di Giosia Jahangir Habib Lea Kenigsberg
Dylanger Pittman Weitao Zhu

Abstract. The Euclidean log convex density theorem, proved by Gregory Chambers in 2015, asserts that in Euclidean space with a log convex density spheres about the origin are isoperimetric. We provide a partial extension to hyperbolic space in which volume and perimeter densities are related but different.

Acknowledgements: We would like to thank the NSF REU grant program, Williams College (including the Finnerty Fund), Stony Brook University, and the Mathematical Association of America for funding. Additionally, we would like to thank our advisor, Frank Morgan, for valuable input and discussions throughout our research.

1 Introduction

The Euclidean log convex density theorem, proved by Gregory Chambers [Ch] in 2015, asserts that in Euclidean space with a log convex density spheres about the origin are isoperimetric (minimize perimeter for given volume). The analogous conjecture in hyperbolic space, \mathbb{H}^n , says that for a log convex density, spheres about the origin are isoperimetric. We prove an easier version in which the perimeter is weighted by a larger density than the volume:

Theorem 7. *Consider \mathbb{H}^n with smooth, radial, log convex volume density $\phi(R)$ and perimeter density*

$$\phi(R) \cdot 2 \cosh^2(R/2),$$

where R is distance from the origin. Then spheres about the origin are uniquely isoperimetric.

Note that a smooth, radial density is log convex if and only if it is a log convex function of R .

The problem of perimeter minimization is an ancient one. Two thousand years ago, Zenodorus showed that planar circles are perimeter minimizers for given enclosed area. Similar results were proven over the past 150 years for spheres in \mathbb{R}^n , \mathbb{S}^n , and \mathbb{H}^n (see Morgan, [Mo, §13.2]). Here, \mathbb{S}^n denotes the n -dimensional unit sphere.

More recently, mathematicians have studied manifolds with density. A density f is a positive function defined on a manifold. For a region Ω , the weighted volume of Ω with respect to density f is

$$\int_{\Omega} f dV,$$

and the weighted area of its boundary is

$$\int_{\partial\Omega} f dA.$$

The interest in manifolds with density is partially due to their role in Perelman's 2002 proof of the Poincaré conjecture [Mo, Chapter 18]. The notion of minimizing weighted area with respect to weighted volume is the underlying principle found in Theorem 7. We break its proof into two steps.

Firstly, we use the Poincaré ball model of hyperbolic space to reduce the problem to the open unit ball in \mathbb{R}^n with equal volume and perimeter densities which diverge at the boundary.

Secondly, Proposition 6 shows that a component of an isoperimetric region must be bounded. Otherwise, we carefully truncate the region and restore the volume elsewhere to get a good candidate which nonetheless must have no less perimeter. This inequality, paired with other differential inequalities relating the boundary of the truncation to the rate of volume growth, forces the volume outside the ball of radius r to reach 0 in finite time.

Structurally, the outline of this paper is as follows: Section 2 provides notation. Section 3 begins with an explicit formula (Lemma 1) relating Poincaré and exponential coordinates

on \mathbb{H}^n . Proposition 3 shows that \mathbb{H}^n featuring two densities is equivalent to the Euclidean ball featuring only one density. Propositions 5 and 6 after Morgan-Pratelli [MP, Thms. 3.3, 5.9] establish the existence and boundedness of isoperimetric regions. The local analysis of Chambers [Ch] for a single density on Euclidean space completes the proof of our main Theorem 7.

2 Notation

Here we provide a reference for the notation used in many of our lemmas and propositions.




Figure 1: By relating the slice E_r to the rate of growth of the volume of E inside the sphere, one can obtain inequalities useful in proving existence and boundedness of isoperimetric regions E .

As in Figure 1, in both hyperbolic and Euclidean space, denote the sphere and closed ball of radius r by $S(r)$ and $B(r)$. Let $|\Omega|$ denote the unweighted measure of a surface or region Ω . For a region E , let E_r be the slice of E by the sphere $S(r)$ (see Morgan, [Mo, 4.11]). Let F_r be the restriction of the boundary of E to the exterior of the ball $B(r)$ and define $P(r)$ as $|F_r|$. Similarly, let $V(r)$ be the volume of the restriction of E to the exterior of the ball $B(r)$. Let $p(r)$ denote the perimeter of E_r . In the presence of any density f , apply f as a subscript to indicate weighted volume or perimeter.

3 Proof of Theorem

After two lemmas, Proposition 3 equates hyperbolic space featuring two densities with the Euclidean ball featuring one density. Propositions 5 and 6 prove existence and boundedness of isoperimetric regions. Our main Theorem 7 now can apply Chambers [Ch] to prove spheres are uniquely isoperimetric in \mathbb{H}^n with our unequal densities.

Lemma 1. *On the Poincaré ball model of \mathbb{H}^n with radial coordinate r , distance from the origin is $2 \tanh^{-1} r$.*

Proof. This distance may be written as

$$\begin{aligned} \int_0^r \frac{2}{1-x^2} dx &= \int_0^r \frac{1}{1-x} + \frac{1}{1+x} dx \\ &= \ln \left(\frac{1+r}{1-r} \right) \\ &= 2 \tanh^{-1} r. \end{aligned}$$

□

A density $\phi(x)$ on the line segment $[0, 1)$ can be converted into a radial density on the open unit ball in \mathbb{R}^n by considering the density $\phi(r)$, where r is Euclidean distance from the origin. Thus, by Lemma 1, ϕ may be similarly converted into a radial density on hyperbolic space as a function of hyperbolic distance from the origin, R , by computing the composition $\phi \circ \tanh(R/2)$. This allows for conversion between radial densities on hyperbolic space and radial densities on the Euclidean ball. Lemma 2 establishes that log convexity of $\phi \circ \tanh(R/2)$ on $[0, \infty)$ implies log convexity of ϕ on $[0, 1)$.

Lemma 2. *Let g be a smooth function with g' positive and g'' negative. If the composition of a smooth positive nondecreasing function f with g is log convex, then f is log convex.*

Proof. By assumption,

$$\begin{aligned} 0 &\leq f^2 \cdot (\log(f \circ g))'' = (g')^2 \cdot (ff'' - (f')^2) + ff'g'' \\ &\leq (g')^2 \cdot (ff'' - (f')^2) = (g' \cdot f)^2 (\log f)'' . \end{aligned}$$

Therefore f is log convex. □

Take any smooth radial density on hyperbolic space. Let $\phi(r)$ be the density as a function of Poincaré radial coordinate r . Then if the density is log convex, ϕ must be nondecreasing. This is because the symmetry and smoothness of the density at the origin implies we have $\phi'(0) = 0$ and log convexity of the density implies convexity of the density, so the quantity ϕ' could not become negative anywhere on $(0, 1)$. By hypothesis, the composition of ϕ with $\tanh(R/2)$ is log convex on $[0, \infty)$. Taking g to be $\tanh(R/2)$ and f to be ϕ provides that ϕ is log convex on $[0, 1)$.

The hyperbolic metric in Poincaré coordinates is a conformal rescaling of that on the Euclidean ball by a factor of $2/(1-r^2)$. By the effects of conformally rescaling a metric, we see that hyperbolic space has the same isoperimetric profile in Poincaré coordinates as the open unit Euclidean ball with with area density $(2/(1-r^2))^{n-1}$ and volume density $(2/(1-r^2))^n$. We are now able to establish the equivalence between hyperbolic space with multiple densities and the Euclidean ball with a single density.

Proposition 3. *The space \mathbb{H}^n with radial volume density $\phi(R)$ and perimeter density*

$$\phi(R) \cdot 2 \cosh^2(R/2)$$

is equivalent to the open unit Euclidean ball with density

$$\phi(2 \tanh^{-1} r) \cdot \left(\frac{2}{1-r^2} \right)^n,$$

where R is hyperbolic distance from the origin and r is Euclidean distance from the origin.

Proof. Note that $\cosh^2(R/2) = 1/(1-\tanh^2(R/2))$. So, the perimeter density may be written as

$$\phi(R) \cdot \frac{2}{1-\tanh^2(R/2)}.$$

By Lemma 1, we may rewrite these functions in Poincaré coordinates to see that the volume density becomes $\phi(2 \tanh^{-1} r)$ and the perimeter density becomes

$$\phi(2 \tanh^{-1} r) \cdot \frac{2}{1-r^2},$$

where r is the radial Poincaré coordinate. Finally, let dV_0 and dA_0 denote the Euclidean volume and area elements. Then in Poincaré coordinates, the weighted volume element is

$$\phi(2 \tanh^{-1} r) \cdot \left(\frac{2}{1-r^2} \right)^n dV_0,$$

and the weighted area element is

$$\phi(2 \tanh^{-1} r) \cdot \left(\frac{2}{1-r^2} \right)^n dA_0.$$

These volume and area elements are equivalent to those on the described weighted Euclidean ball. □

Having demonstrated this equivalence between \mathbb{H}^n and the unit ball with certain densities, we are now ready to make statements about existence and boundedness of isoperimetric regions. First, we need a brief lemma relating area of slices of a region to the perimeter of its boundary.

Lemma 4. *Let E be a region in either \mathbb{H}^n or the unit ball in \mathbb{R}^n with continuous, nondecreasing radial volume density, with infinite weighted radial distance to the boundary in the ball case. If E has finite weighted volume then for any r ,*

$$P(r) \geq |E_r|.$$

Proof. Since the radial projection π_r of the exterior F_r onto the sphere $S(r)$ is area nonincreasing, it suffices to show that $\pi_r(F_r)$ covers E_r , up to set of measure zero. If not, then a set of positive measure in E_r is not in the projection. The product of this set with either (r, ∞) or $(r, 1)$, depending on the space, is contained in E . But this set must have infinite weighted volume, violating our assumption of finite weighted volume. \square

Now we may show the existence of isoperimetric regions in our space. Note that this proof follows a similar structure of an existence proof in Morgan-Pratelli [MP, Thm. 3.3].

Proposition 5. *Isoperimetric regions exist in \mathbb{H}^n with nondecreasing, continuous radial volume density $g(R)$ and radial perimeter density $g(R) \cdot 2 \cosh^2(R/2)$, where R is distance from the origin, for all volumes.*

Proof. By Proposition 3, it is enough to show that isoperimetric regions exist on the open unit Euclidean ball with radial density

$$f(r) = g(2 \tanh^{-1} r) \cdot \left(\frac{2}{1-r^2} \right)^n.$$

Consider a sequence of regions of the prescribed volume with perimeter tending to the infimum. By compactness [Mo, §9.1] we may assume convergence of a subsequence to a perimeter-minimizing region.

The difficulty is that the enclosed volume may be strictly less than the prescribed volume, that some volume disappears to infinity. In that case, for some $\epsilon > 0$, for all r , for a tail of the sequence, the volume outside the ball of radius r about the origin is at least ϵ . Fix R and such a region E , determined by R , in the tail of the sequence. We have

$$\int_R^1 |E_r| f(r) dr \geq \epsilon.$$

Because f satisfies the conditions of Lemma 4, we know $|E_r|$ converges to zero as r tends to 1, as $|E_r|$ is bounded above by $P(r)$ which must also converge to 0. Hence, when r is chosen close enough to 1, $|E_r|$ is no larger than half the unweighted surface area of the sphere of radius r . When this condition holds, we may apply the standard isoperimetric inequality,

$$p(r) \geq c |E_r|^{\frac{n-2}{n-1}},$$

where c is some dimensional constant.

Let M be the supremum of $|E_r|$ for $r \geq R$, which is finite by Lemma 4. In addition, Lemma 4 implies that

$$|\partial E|_f \geq M f(R). \tag{1}$$

To see why this relationship holds, let $\{E_{r_n}\}$ be a sequence of slices with area tending to supremum M , with $r_n \geq R$. Then we have that

$$\begin{aligned} P(r_n) &\geq |E_{r_n}|, \\ P_f(r_n) &\geq |E_{r_n}|f(r_n), \\ |\partial E|_f &\geq |E_{r_n}|f(R), \end{aligned}$$

where we have used that f is nondecreasing. Taking a limit produces the desired inequality.

Take note that the function $P(r)$ is nonincreasing, so it must be differentiable almost everywhere, and by extension, $P_f(r)$ is differentiable almost everywhere. By standard slicing theory [Mo, §4.11], for almost all r ,

$$f(r)p(r) \leq -P'_f(r).$$

Since any singular changes in P_f are positive, integration yields that for R close enough to 1, we have

$$\begin{aligned} |\partial E|_f &\geq \int_R^1 -P'_f(r) dr \geq \int_R^1 p(r)f(r)dr \\ &\geq c \int_R^1 |E_r|^{\frac{n-2}{n-1}} f(r)dr \geq \frac{c}{M^{\frac{1}{n-1}}} \int_R^1 |E_r|f(r)dr \\ &\geq \frac{c}{M^{\frac{1}{n-1}}} \epsilon. \end{aligned}$$

By (1),

$$|\partial E|_f^{\frac{n}{n-1}} \geq cf(R)^{\frac{1}{n-1}} \epsilon. \quad (2)$$

Note that because the weighted perimeters of the regions in the sequence converge, the sequence of perimeters is bounded by some positive k . Because E is one of the sequence elements, (2) provides that

$$\frac{\epsilon k^{\frac{n}{n-1}}}{c} \geq f(R)^{\frac{1}{n-1}}. \quad (3)$$

This implies that f is bounded above. Recalling the definition of f and noting that g is bounded below provides that f must be unbounded, a contradiction.

Therefore there is no loss of volume to infinity and the limit provides the desired perimeter-minimizing region. \square

Proposition 6 establishes that these isoperimetric regions must be bounded. See Morgan-Pratelli [MP, Thm. 5.9] for a similar proof in \mathbb{R}^n .

Proposition 6. *Isoperimetric regions are bounded in \mathbb{H}^n with nondecreasing, continuous radial volume density $g(R)$ and radial perimeter density $g(R) \cdot 2 \cosh^2(R/2)$, where R is distance from the origin.*

Proof. Denote the volume and perimeter densities by g and f .

Suppose we have such an isoperimetric region E with $V_g(R)$ never zero. Because the volume density is continuous and nondecreasing, and because E encloses finite weighted volume, Lemma 4 implies that

$$|E_r| \leq P(r).$$

Because f is nondecreasing,

$$|E_r|_f \leq P_f(r).$$

The standard isoperimetric inequality on the sphere tells us that for any r for which $|E_r|$ is at most half of the area of $S(r)$, one has

$$p(r) \geq c|E_r|^{\frac{n-2}{n-1}}, \quad (4)$$

where c is some positive dimensional constant. Because $P(r)$ is bounded by $|\partial E|$, which is finite because $|\partial E|_f$ is finite and f is bounded below, for all r large enough $|E_r|$ is at most half of the area of $S(r)$ and (4) holds. Because the f density on E_r is constant, we may multiply inequality (4) by $f(r) = g(r) \cdot 2 \cosh^2(r/2)$ to obtain, for all r large enough,

$$p_f(r) \geq c|E_r|_f^{\frac{n-2}{n-1}} g(r)^{\frac{1}{n-1}} \cosh^{\frac{2}{n-1}}(r/2),$$

where we've absorbed the power of 2 into constant c . Noting that g is bounded below by some constant we may absorb this lower bound into a new constant c to obtain

$$\begin{aligned} p_f(r) &\geq c|E_r|_f^{\frac{n-2}{n-1}} \cosh^{\frac{2}{n-1}}(r/2), \\ &\geq c|E_r|_f \cdot |E_r|_f^{\frac{-1}{n-1}} \cosh^{\frac{2}{n-1}}(r/2), \\ &\geq c|E_r|_f \cdot P_f(r)^{\frac{-1}{n-1}} \cosh^{\frac{2}{n-1}}(r/2). \end{aligned} \quad (5)$$

Note that we have taken negative powers of $|E_r|$, which may be zero if E is disconnected. However, for such values of r , every quantity in the inequality is zero and so the statements hold trivially. Furthermore, we are justified in taking a negative power of $P_f(r)$ as it is never zero, because E is unbounded.

Because $V(r)$, $P(r)$, and f are monotonic functions, $V_f(r)$ and $P_f(r)$ are differentiable almost everywhere. By Morgan [Mo, §4.11], the differential relationships (6) and (7) hold almost everywhere:

$$-P'_f(r) \geq p_f(r) \quad (6)$$

$$-2 \cosh^2(r/2) V'_g(r) = -V'_f(r) = |E_r|_f. \quad (7)$$

Hence, for almost all r large enough, (5) may be rewritten using (6) and (7) as

$$\begin{aligned}
 -P'_f(r) &\geq c(-V'_g(r))P_f(r)^{\frac{-1}{n-1}} \cosh^{\frac{2}{n-1}}(r/2), \\
 -P'_f(r)P_f(r)^{\frac{1}{n-1}} &\geq -2 \cdot cV'_g(r) \cosh^{\frac{2n}{n-1}}(r/2), \\
 -(P_f^{\frac{n}{n-1}})'(r) &\geq -cV'_g(r) \cosh^{\frac{2n}{n-1}}(r/2),
 \end{aligned}$$

where c may change from line to line. Because this inequality holds for almost all r large enough, we may perform an integration from all large r to ∞ . The quantity $-(P_f^{\frac{n}{n-1}})$ may not be continuous, but it is increasing, so the integral of its derivative is no larger than the absolute change in $P_f^{\frac{n}{n-1}}$. Knowing that $V_g(r)$ is continuous provides that the integral of its derivative is exactly the change in $V_g(r)$. Finally, noting that $P_f(r)$ and $V_g(r)$ converge to 0 at ∞ , we integrate starting at an r large enough,

$$\begin{aligned}
 P_f^{\frac{n}{n-1}}(r) &\geq \int_r^\infty -(P_f^{\frac{n}{n-1}})'(x) dx, \\
 P_f^{\frac{n}{n-1}}(r) &\geq c \cosh^{\frac{2n}{n-1}}(r/2) \int_r^\infty -V'_g(x) dx, \\
 P_f(r) &\geq c \cosh^2(r/2)(V_g(r))^{\frac{n-1}{n}}, \tag{8}
 \end{aligned}$$

where the positive constant c may be modified from line to line.

The goal in the remainder of the proof is to derive a contradiction by shifting small pieces of volume of E from far away to places closer to the center. We would like to do this in a way that does not increase the perimeter of the isoperimetric region too dramatically.

On this note, for large R , there is a small positive ϵ and an inflated version of E , E_ϵ , that agrees with E outside $B(R)$ and contains ϵ more g -weighted volume satisfying

$$|\partial E_\epsilon|_f \leq |\partial E|_f + \epsilon(H + 1),$$

where H denotes the constant generalized unaveraged mean curvature of ∂E with respect to densities f and g . To understand why this version of E exists, recall that H is the derivative of f -weighted perimeter with respect to small increases of g -weighted volume. So, there is a small number $\bar{\epsilon}$ such that $\epsilon \leq \bar{\epsilon}$ implies

$$\Delta P \leq \epsilon(H + 1),$$

where ΔP is understood as the increase in f -weighted perimeter of ∂E from adding g -weighted volume ϵ to E somewhere relatively close to the origin. Noting that $V_g(r)$ converges to 0 at ∞ , we may pick R large enough to ensure both that $V_g(R) = \epsilon \leq \bar{\epsilon}$ and that the necessary inflation of E by ϵ units of g -weighted volume happens within $B(R)$. Finally, once ϵ is determined by R , we can construct the described E_ϵ by inflating E inside $B(R)$ with ϵ units of g -weighted volume.

Fix R large, as determined by $\bar{\epsilon}$ and the requirements in (8). Let \bar{E} denote the restriction of E_ϵ to the ball $B(R)$ and note that this truncation provides $|E|_g = |\bar{E}|_g$. By (8),

$$\begin{aligned}
|\partial\bar{E}|_f &= |\partial E_\epsilon|_f - P_f(R) + |E_R|_f \\
&\leq |\partial E|_f + \epsilon(H+1) - c \cosh^2(R/2)\epsilon^{\frac{n-1}{n}} \\
&\quad + |E_R|_f.
\end{aligned} \tag{9}$$

Since E is an isoperimetric region, $|\partial\bar{E}|_f \geq |\partial E|_f$, as they enclose the same volume, and (9) becomes

$$0 \leq \epsilon(H+1) - c \cosh^2(R/2)\epsilon^{\frac{n-1}{n}} + |E_R|_f.$$

Recall that ϵ is determined by R and can be made arbitrarily small by making R arbitrarily large. For ϵ very small, the linear term is dwarfed by the power term. Hence, for any R large enough, modifying the constant c as necessary,

$$\begin{aligned}
c \cosh^2(R/2)\epsilon^{\frac{n-1}{n}} &\leq |E_R|_f = -2 \cosh^2(R/2)V'_g(R), \\
c(V_g(R))^{\frac{n-1}{n}} &\leq -V'_g(R), \\
c &\leq -(V_g^{\frac{1}{n}})'(R).
\end{aligned}$$

This last estimate gives a contradiction with the assumption $V_g(r) > 0$ for all r , and the proof is complete. \square

Now that we know by Propositions 5 and 6 that isoperimetric regions exist and are bounded, we can apply the local analysis in Chambers to conclude they are balls about the origin.

Theorem 7. *Consider \mathbb{H}^n with smooth, radial, log convex volume density $\phi(R)$ and perimeter density*

$$\phi(R) \cdot 2 \cosh^2(R/2),$$

where R is distance from the origin. Then spheres about the origin are uniquely isoperimetric.

Proof. By Proposition 3, we may reduce the problem to the open unit Euclidean ball with volume and perimeter density $\phi(2 \tanh^{-1}(r)) \cdot (2/(1-r^2))^n$. By Lemma 2, this is a log convex density on the open unit ball. Because there is infinite weighted distance to the boundary, Proposition 5 guarantees that isoperimetric regions exist. These isoperimetric regions are bounded by Proposition 6.

Because this isoperimetric region is bounded in a subset of \mathbb{R}^n with smooth, log convex, radial density, we may apply the same local analysis done by Chambers [Ch] on isoperimetric regions in \mathbb{R}^n . From this, we conclude that spheres about the origin are isoperimetric, uniquely because the density is strictly log convex. \square

Remark. The proof of our main Theorem 7 compared hyperbolic space featuring perimeter density much larger than volume density with Euclidean space featuring equal densities. Since the log convex density conjecture for \mathbb{H}^n involves equal densities, the comparison would be with the Euclidean ball featuring perimeter density much smaller than volume density, about which little is known.

Conjecture 8. (*The log convex density conjecture on hyperbolic space*). In \mathbb{H}^n with smooth, log convex radial density, every sphere about the origin is isoperimetric.

References

[Ch] Gregory Chambers. Proof of the log convex density conjecture. J. Eur. Math. Soc., to appear.

[Mo] Frank Morgan. *Geometric Measure Theory*. 5th ed., 2016.

[MP] Frank Morgan and Aldo Pratelli. Existence of isoperimetric regions in \mathbb{R}^n with density, *Ann. Global Anal. Geom.* 43 (2013), 331-365.